

## The effect of resveratrol on expression of matrix metalloproteinase 9 and its tissue inhibitors in vascular smooth muscle cells

Effat Farrokhi<sup>(1)</sup> , Keihan Ghatreh-Samani<sup>(2)</sup> , Najmeh Salehi-Vanani<sup>(3)</sup>, Amin Mahmoodi<sup>(3)</sup>

### Original Article

#### Abstract

**BACKGROUND:** Matrix metalloproteinase 9 (MMP-9) is involved in extracellular matrix degradation and remodeling. An increase in MMP-9 expression by vascular component cells plays an important role in atherosclerotic plaque formation and rupture. Resveratrol, a polyphenolic substance, was suggested to play a role in preventing the progress of atherosclerotic disease. The aim of this study was to investigate the effect of resveratrol on MMP-9 and tissue inhibitors of metalloproteinases (TIMPs) in vascular smooth muscle cells (VSMCs) after treatment with H<sub>2</sub>O<sub>2</sub>.

**METHODS:** Cultured VSMCs were pre-treated with 0.2 mM of H<sub>2</sub>O<sub>2</sub> before stimulation with different concentration of resveratrol. Expression of MMP-9, TIMP-1, and TIMP-3 genes were measured using real-time polymerase chain reaction (PCR) method, and MMP-9 protein level was detected using western blot analysis.

**RESULTS:** Resveratrol at 120 µmol/l concentration reduced the elevated level of MMP-9 induced by H<sub>2</sub>O<sub>2</sub> in VSMCs as  $1.85 \pm 0.35$  folds ( $P < 0.050$ ) and  $8.70 \pm 1.20$  folds ( $P < 0.050$ ) after 24 and 48 hours, respectively. Resveratrol increased the diminished level of TIMP-1 induced by H<sub>2</sub>O<sub>2</sub> as  $2.5 \pm 0.48$  folds following the treatment with 120 µmol/l after 48 hours ( $P < 0.050$ ).

**CONCLUSION:** Resveratrol as an antioxidant can decrease MMP-9 production, not only by suppressing MMP-9 expression, but also by augmenting TIMP-1 production. Altogether, resveratrol as an antioxidant can regulate the MMP-9/TIMP-1 balance, and may be considered as a preservative agent in the treatment and prevention of atherosclerosis.

**Keywords:** Matrix Metalloproteinase 9, TIMPS, Resveratrol, Vascular Smooth Muscle

*Date of submission:* 29 Aug. 2016, *Date of acceptance:* 21 May 2018

#### Introduction

Atherosclerosis is the most common cause of death in the world, and oxidative stress has a critical role in the pathogenesis and development of atherosclerosis. Matrix metalloproteinase 9 (MMP-9) is a member of the family of endopeptidases, that is involved in extracellular matrix degradation and remodeling. Increased expression of MMP-9 by vascular component cells during vascular injury and inflammation plays an important role in atherosclerotic plaque formation and rupture.<sup>1</sup>

MMPs activity is regulated by tissue inhibitors of metalloproteinases (TIMPs).<sup>2</sup> Changes in TIMP levels are considered to be important due to their direct effect on MMP activity. Resveratrol is a natural polyphenolic antioxidant, and several studies

have confirmed its protective effects on the cardiovascular system in patients with coronary artery disease (CAD).<sup>3,4</sup>

Resveratrol has a cardioprotective activity via various mechanisms such as anti-inflammatory and antioxidant activity.<sup>5,6</sup>

Resveratrol reduces neural damage after cerebral ischemia. The elevated levels of MMP-9 were notably diminished in the resveratrol-treated mice as compared to the vehicle MCAO mice; suggesting that resveratrol can protect neural tissue against acute ischemic stroke, which could be attributed to its activity against MMP-9 by inhibiting JNK and PKC signal transduction.<sup>7,8</sup>

It seems that inhibition of MMPs activation or prevention of their upregulation could inhibit the

1- Assistant Professor, Department of Molecular Medicine, School of Advanced Technologies, Shahrekord University of Medical Sciences, Shahrekord, Iran

2- Associate Professor, Cellular and Molecular Research Center, Basic Health Sciences Institute, Shahrekord University of Medical Sciences, Shahrekord, Iran

3- Clinical Biochemistry Research Center, Basic Health Sciences Institute, Shahrekord University of Medical Sciences, Shahrekord, Iran  
Correspondence to: Keihan Ghatreh-Samani, Email: [ghatreh\\_k@skums.ac.ir](mailto:ghatreh_k@skums.ac.ir)

effects of oxidative stress on atherosclerotic plaque formation. The aim of this study was to investigate the effect of resveratrol on MMP-9 and TIMPs in vascular smooth muscle cells (VSMCs) after treatment with H<sub>2</sub>O<sub>2</sub>.

### Materials and Methods

In this experimental study, human aorta-vascular smooth muscle cells (HA-VSMC) were purchased from cell bank of Pasteur Institute of Iran in Tehran, Iran. The cells were cultured in F12K growth medium. F12K media contained 10% fetal bovine serum (FBS), 0.05 mg/ml ascorbic acid, 0.01 mg/ml insulin, 0.05 mg/ml transferrin, 10 ng/ml sodium selenite, 0.03 mg/ml human epidermal growth factor (EGFs), 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) to a final concentration of 10 mM, penicillin (100 U/ml), and streptomycin (100 U/ml).

Cells were cultured in a humidified incubator containing 5% CO<sub>2</sub> at 37 °C. Daily control of cell growth and cell division in culture condition were done. Cells used for the experiments were 3 to 7 passages.

3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay was used to determine resveratrol concentration causing 50% cell death (the half maximal inhibitory concentration or IC<sub>50</sub>). Briefly, the cells were seeded at a density of  $15 \times 10^3$  in 96-well plates. After achieving approximately 80% confluence, cells were treated with increasing concentration of resveratrol (40 to 200 µmol/l) for 48 hours. Thereafter, 20 µl MTT (5 mg/ml, Sigma-Aldrich, Germany) solution was added to each well. The formazan crystal was dissolved by adding 150 µl dimethyl sulfoxide (DMSO) to each well after 4 hours, and the optical density of each well was

measured at 490 nm on an enzyme-linked immunosorbent assay (ELISA) plate reader (Awareness, USA) following 4 hours of incubation at 37 °C.

The cells were seeded in a 12-well plate at a density of  $1 \times 10^4$  cells per well. When the cells achieved approximately 80% confluence, they were pretreated with physiologic concentration (0.2 mM) of H<sub>2</sub>O<sub>2</sub> for inducing oxidative stress.<sup>9</sup> Then, the cells co-cultured with 80, 100, and 120 µmol/l resveratrol (Sigma, USA) for 24 and 48 hours.

**RNA isolation and cDNA synthesis:** Total RNA was extracted from the cells using Trizol (Invitrogen, USA) according to the manufacturer instructions. RNA was quantified using a Nanodrop 2000C spectrophotometer (Thermo Scientific, USA) and treated with DNase. Then, cDNA was synthesized from 0.5 mg total RNA using random primer and the cDNA Synthesis kit (Thermo Fisher Scientific, USA).

**Real-time polymerase chain reaction (Real-time PCR):** The expression of MMP-9, TIMP-1, and TIMP-3 genes was measured using quantitative real-time PCR. The experiments were performed using Rotor-Gene 3000 real-time DNA amplification system (Corbett Research, Australia) and SYBR green method. Primers used for real-time PCR are listed in table 1. Experiments were performed in triplicate,<sup>6</sup> using 5 µl SYBR green PCR Master Mix, 0.2 µl primer sets, 2 µl cDNA (40 ng), and 3.6 µl nuclease-free H<sub>2</sub>O to yield a 10 µl reaction. The amplification conditions were as follow: initial denaturing at 94 °C for 5 minutes, then 40 cycles of 95 °C for 15 seconds, 59 °C for 20 seconds, and 72 °C for 30 seconds. The comparative cycle threshold (CT) ( $\Delta\Delta CT$ ) method was used for data quantitation, and glyceraldehyde-3-phosphate dehydrogenase (GAPDH) gene expression was used as an endogenous reference.

**Table 1.** Primer sequences and product length

Genes	Genes Primer sequences (5'-3')	Amplicon length (bp)	Gene bank reference
MMP-9	Forward: GCTCACCTTCACTCGCGTGTA Reverse: TCCGTGCTCCGCGACA	70	NM_004994.2
TIMP-1	Forward: CCTGGCTTCTGGCATCCTG Reverse: CCACGAACCTGGCCCTGATG	125	NM_003254.2
TIMP-3	Forward: GTCGCGTCTATGATGGCAAG Reverse: AAGCAAGGCAGGTAGTAGCA	151	NM_000362.4
GAPDH	Forward: ACACCCACTCCTCCACCTTTG Reverse: TCCACCACCCTGTTGCTGTAG	112	NM-002046.5

MMP-9: Matrix metalloproteinase 9;TIMP: Tissue inhibitors of metalloproteinases; GAPDH: Glyceraldehyde-3-phosphate dehydrogenase

**Western blot analysis:** Western blot analysis was used for detecting MMP-9 protein expression. VSMCs were lysed in ice-cold radioimmune precipitation assay buffer (6 ×) containing protease inhibitor cocktail. Samples were electrophoresed on 12% sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE). The separated proteins were transferred to nitrocellulose membranes, and incubated overnight at 4 °C with blocking solution [5% nonfat dried milk in phosphate-buffered saline (PBS) containing 0.1% Tween 20]. Membranes were incubated with MMP-9 antibody (1:3000) at room temperature for 2 hours. After washing, the membrane was incubated with goat anti-rabbit IgG horseradish peroxidase conjugate (1:10,000) at room temperature for 90 minutes. Finally, the color was developed with the addition of 3,3',5,5'-tetramethylbenzidine membrane peroxidase substrate. Beta-actin was detected as a sample loading control.

All experiments were done in triplicate. Results are expressed as the mean ± standard error of mean (SEM). Statistical analysis was done using nonparametric Kruskal-Wallis test. The nonparametric Mann-Whitney U test was used to compare differences between control and test groups. P value of less than 0.050 was considered as the level of significance.

## Results


The IC<sub>50</sub> of resveratrol estimated by MTT assay was about 120 µmol/l. This concentration of resveratrol was chosen for further experiments.

In this study, the resveratrol effects on expression of MMP-9 and its inhibitors (TIMP-1 and -3) were investigated in VSMCs after inducing with H<sub>2</sub>O<sub>2</sub>. H<sub>2</sub>O<sub>2</sub> used at the non-toxic concentration of 0.2 mM. MMP-9 expression increased  $1.43 \pm 0.29$  and  $1.98 \pm 0.54$  folds after 24 and 48 hours, respectively ( $P < 0.050$  for both) after treatment of the cells with H<sub>2</sub>O<sub>2</sub> without resveratrol. In contrast, resveratrol at different concentrations decreased MMP-9 expression, when given simultaneously with H<sub>2</sub>O<sub>2</sub>.

After 24 hours, MMP-9 expression was decreased  $1.60 \pm 0.21$ ,  $1.57 \pm 0.30$ , and  $1.85 \pm 0.35$  folds following the treatment with 80, 100, and 120 µmol/l resveratrol when compared with the H<sub>2</sub>O<sub>2</sub>-treated group ( $P < 0.050$  for all) (Figure 1).

After 48 hours, resveratrol at 80, 100, and 120 µmol/l concentrations reduced the elevated level of MMP-9 induced by H<sub>2</sub>O<sub>2</sub> as  $6.20 \pm 1.28$ ,  $5.50 \pm 1.96$ , and  $8.70 \pm 1.20$  folds, respectively,

when compared with the H<sub>2</sub>O<sub>2</sub>-treated group ( $P < 0.050$  for all) (Figure 1).




**Figure 1.** H<sub>2</sub>O<sub>2</sub> upregulated the expression of matrix metalloproteinase 9 (MMP-9) in vascular smooth muscle cells (VSMCs). Resveratrol at 80, 100, and 120 µmol/l concentrations reduced the elevated level of MMP-9 induced by H<sub>2</sub>O<sub>2</sub>.

\*  $P < 0.050$  compared with control group

#  $P < 0.050$  compared with H<sub>2</sub>O<sub>2</sub>-treated group

Western blot analysis confirmed the changes observed at MMP-9 mRNA level (Figure 2). In western blot analysis, beta-actin (42 kDa) was used as internal control.




**Figure 2.** The expression of matrix metalloproteinase protein 9 (MMP-9) determined by western blot analysis after treatment with 80, 100, 120 µmol/l resveratrol in vascular smooth muscle cells (VSMCs). Beta-actin (42 kDa) was used as an internal control to standardize the protein loading in western blotting.

Lane 1: control; Lane 2: treated with H<sub>2</sub>O<sub>2</sub>; Lane 3-5: treated with H<sub>2</sub>O<sub>2</sub> and various concentration of resveratrol (80, 100, and 120 µmol/l, respectively).

After treating the cells with H<sub>2</sub>O<sub>2</sub> without resveratrol, TIMP-1 expression decreased  $1.73 \pm 0.26$  folds after 48 hours ( $P < 0.050$ ). In contrast, resveratrol at different concentrations increased TIMP-1 expression when given simultaneously with H<sub>2</sub>O<sub>2</sub>.

After 48 hours, resveratrol at 80, 100, and 120 µmol/l concentrations increased the diminished level of TIMP-1 induced by H<sub>2</sub>O<sub>2</sub> about  $1.74 \pm 0.51$ ,

1.93  $\pm$  0.37, and 2.50  $\pm$  0.48 folds, respectively, compared with the H<sub>2</sub>O<sub>2</sub>-treated group ( $P < 0.050$  for all) (Figure 3). There was no significant change in the expression of TIMP-1 after 24 hours.



**Figure 3.** H<sub>2</sub>O<sub>2</sub> downregulated the expression of tissue inhibitors of metalloproteinase 1 (TIMP-1) in vascular smooth muscle cells (VSMCs). Resveratrol in 80, 100, and 120  $\mu$ mol/l concentrations increased the diminished level of TIMP-1 induced by H<sub>2</sub>O<sub>2</sub> after 48 hours.

\*  $P < 0.050$  compared with control group

#  $P < 0.050$  compared with H<sub>2</sub>O<sub>2</sub>-treated group

We found no significant effect by resveratrol on TIMP-3 expression by real time PCR (not shown).

## Discussion

In this study, we investigated the resveratrol effect on MMP-9 expression in H<sub>2</sub>O<sub>2</sub>-induced VSMCs. Resveratrol decreased MMP-9 expression in H<sub>2</sub>O<sub>2</sub>-induced VSMCs.

Matrix metalloproteinases play important roles in cardiovascular diseases, and oxidative stress has destructive effect on vascular biology through the activation of MMPs.<sup>10</sup>

MMP-9 is involved in the breakdown of extracellular matrix proteins in various physiological and pathological conditions, such as angiogenesis, bone development, and atherosclerosis plaque rupture.<sup>1</sup>

Enhanced expression and activities of MMP-9 have been observed in vascular injury. TIMP-1 is endogenous inhibitor of MMP-9, and it is suggested that TIMP-1 plays an important role in cardiovascular disease.

Regarding the effects of oxidative stress in vascular remodeling, antioxidant approaches are used to reduce the upregulation of MMPs, and attenuate the tissue dysfunction and remodeling during vascular diseases.<sup>11-13</sup>

In this study, H<sub>2</sub>O<sub>2</sub> increased MMP-9 expression in VSMCs. Upregulation of MMP-9 is not unexpected in conditions that free radicals were increased by H<sub>2</sub>O<sub>2</sub> or other oxidants. Elevated levels of MMP-9 due to oxidant stress in VSMCs has been demonstrated in many studies.<sup>10,14</sup>

Resveratrol decreased MMP-9 expression in H<sub>2</sub>O<sub>2</sub>-induced VSMCs. Resveratrol displays potent antioxidant activity, thereby can scavenge free radicals produced by H<sub>2</sub>O<sub>2</sub>, and so reduce the oxidative effects of H<sub>2</sub>O<sub>2</sub> in the environment.

Previous studies have demonstrated that polyphenols interfere with MMP-9 expression, and reduce inflammatory angiogenesis through MMP-9 inhibition.<sup>15,16</sup> The results of this study and other studies<sup>17,18</sup> support a potential protective role for dietary polyphenols in cardiovascular diseases.

Resveratrol inhibits the nuclear factor kappa-light-chain-enhancer of activated B cells (NF- $\kappa$ B) signal transduction pathway that mediates the expression of MMPs and many other genes involved in inflammatory and pro-oxidant processes associated with vascular disease development.<sup>19</sup> The capability of resveratrol in reducing of NF- $\kappa$ B activity may be considered indicative to change MMP-9 expression in this study.

The MMPs are tightly regulated, not only at the transcriptional level, but also by their specific inhibitors.<sup>20,21</sup> The effect of antioxidant on MMP or TIMP expression in different cells has been previously reported.<sup>22-24</sup>

Epigallocatechin-3-gallate, as an antioxidant agent, reduces the activity and expression of MMP-9, and enhances the expression of TIMP-1 in cancer cell line MDA-MB-231.<sup>22</sup>

In a recent study, resveratrol attenuated blood-brain barrier dysfunction via regulation of MMP-9 and TIMP-1.<sup>23</sup>

According to a study, use of polyphenol extract from olive pomace oil down-regulated the levels of matrix MMP-2 and MMP-9, and increased TIMP-1 expression in human endothelial cells,<sup>24</sup> which could represent a powerful tool for the prevention and treatment of endothelial dysfunction-associated vascular disease.

Matrix metalloproteinases contribute to plaque rupture, atherothrombosis, and myocardial infarction.<sup>25</sup> For this reason, using MMP inhibitors can be considered as a preservative agent for the prevention and treatment of cardiovascular diseases.<sup>26</sup>

The results of this study showed that resveratrol, as an antioxidant, can decrease MMP-9 expression,



not only by suppressing MMP-9 production, but also by augmenting TIMP-1 production.

It seems that due to the dual role of resveratrol with regard to regulation of the MMP-9/TIMP-1 balance, it could be a good choice for preventing and treating cardiovascular diseases.

### Conclusion

Altogether, the results of this study showed that resveratrol, as an antioxidant, can regulate the MMP-9/TIMP-1 balance, and may be considered as a preservative agent in the prevention and treatment of atherosclerosis. To better understand the atherosclerosis prevention and treatment with resveratrol, further studies with animal models are necessary.

### Acknowledgments

This work was obtained from MSc. thesis. Hereby, we gratefully thank Research and Technology Deputy of Shahrekord University of Medical Science, Shahrekord, Iran.

### Conflict of Interests

Authors have no conflict of interests.

### References

- Hobeika MJ, Thompson RW, Muhs BE, Brooks PC, Gagne PJ. Matrix metalloproteinases in peripheral vascular disease. *J Vasc Surg* 2007; 45(4): 849-57.
- Brew K, Dinakarpanian D, Nagase H. Tissue inhibitors of metalloproteinases: Evolution, structure and function. *Biochim Biophys Acta* 2000; 1477(1-2): 267-83.
- Hung LM, Chen JK, Huang SS, Lee RS, Su MJ. Cardioprotective effect of resveratrol, a natural antioxidant derived from grapes. *Cardiovasc Res* 2000; 47(3): 549-55.
- Magyar K, Halmosi R, Palfi A, Feher G, Czopf L, Fulop A, et al. Cardioprotection by resveratrol: A human clinical trial in patients with stable coronary artery disease. *Clin Hemorheol Microcirc* 2012; 50(3): 179-87.
- Birrell MA, McCluskie K, Wong S, Donnelly LE, Barnes PJ, Belvisi MG. Resveratrol, an extract of red wine, inhibits lipopolysaccharide induced airway neutrophilia and inflammatory mediators through an NF-kappaB-independent mechanism. *FASEB J* 2005; 19(7): 840-1.
- Yu W, Fu YC, Zhou XH, Chen CJ, Wang X, Lin RB, et al. Effects of resveratrol on H<sub>2</sub>O<sub>2</sub>-induced apoptosis and expression of SIRT1s in H9c2 cells. *J Cell Biochem* 2009; 107(4): 741-7.
- Gao D, Zhang X, Jiang X, Peng Y, Huang W, Cheng G, et al. Resveratrol reduces the elevated level of MMP-9 induced by cerebral ischemia-reperfusion in mice. *Life Sci* 2006; 78(22): 2564-70.
- Woo JH, Lim JH, Kim YH, Suh SI, Min DS, Chang JS, et al. Resveratrol inhibits phorbol myristate acetate-induced matrix metalloproteinase-9 expression by inhibiting JNK and PKC delta signal transduction. *Oncogene* 2004; 23(10): 1845-53.
- Byon CH, Javed A, Dai Q, Kappes JC, Clemens TL, Darley-Usmar VM, et al. Oxidative stress induces vascular calcification through modulation of the osteogenic transcription factor Runx2 by AKT signaling. *J Biol Chem* 2008; 283(22): 15319-27.
- Chen Q, Jin M, Yang F, Zhu J, Xiao Q, Zhang L. Matrix metalloproteinases: Inflammatory regulators of cell behaviors in vascular formation and remodeling. *Mediators Inflamm* 2013; 2013: 928315.
- Castro MM, Rizzi E, Rodrigues GJ, Ceron CS, Bendhack LM, Gerlach RF, et al. Antioxidant treatment reduces matrix metalloproteinase-2-induced vascular changes in renovascular hypertension. *Free Radic Biol Med* 2009; 46(9): 1298-307.
- Kaneko H, Anzai T, Morisawa M, Kohno T, Nagai T, Anzai A, et al. Resveratrol prevents the development of abdominal aortic aneurysm through attenuation of inflammation, oxidative stress, and neovascularization. *Atherosclerosis* 2011; 217(2): 350-7.
- Saneipour M, Ghatreh-Samani K, Heydari E, Farrokhi E, Abdian N. Adiponectin inhibits oxidized low density lipoprotein-induced increase in matrix metalloproteinase 9 expression in vascular smooth muscle cells. *ARYA Atheroscler* 2015; 11(3): 191-5.
- Rajagopalan S, Meng XP, Ramasamy S, Harrison DG, Galis ZS. Reactive oxygen species produced by macrophage-derived foam cells regulate the activity of vascular matrix metalloproteinases in vitro. Implications for atherosclerotic plaque stability. *J Clin Invest* 1996; 98(11): 2572-9.
- Li YT, Shen F, Liu BH, Cheng GF. Resveratrol inhibits matrix metalloproteinase-9 transcription in U937 cells. *Acta Pharmacol Sin* 2003; 24(11): 1167-71.
- Scoditti E, Calabriso N, Massaro M, Pellegrino M, Storelli C, Martines G, et al. Mediterranean diet polyphenols reduce inflammatory angiogenesis through MMP-9 and COX-2 inhibition in human vascular endothelial cells: A potentially protective mechanism in atherosclerotic vascular disease and cancer. *Arch Biochem Biophys* 2012; 527(2): 81-9.
- Meng L, Liu L, Zhou C, Pan S, Zhai X, Jiang C, et al. Polyphenols and polypeptides in chinese rice wine inhibit homocysteine-induced proliferation

- and migration of vascular smooth muscle cells. *J Cardiovasc Pharmacol* 2016; 67(6): 482-90.
18. Setozaki S, Minakata K, Masumoto H, Hirao S, Yamazaki K, Kuwahara K, et al. Prevention of abdominal aortic aneurysm progression by oral administration of green tea polyphenol in a rat model. *J Vasc Surg* 2017; 65(6): 1803-12.
  19. Ma C, Wang Y, Dong L, Li M, Cai W. Anti-inflammatory effect of resveratrol through the suppression of NF-kappaB and JAK/STAT signaling pathways. *Acta Biochim Biophys Sin (Shanghai)* 2015; 47(3): 207-13.
  20. Woessner JF Jr. Matrix metalloproteinases and their inhibitors in connective tissue remodeling. *FASEB J* 1991; 5(8): 2145-54.
  21. Murphy G, Docherty AJ, Hembry RM, Reynolds JJ. Metalloproteinases and tissue damage. *Br J Rheumatol* 1991; 30(Suppl 1): 25-31.
  22. Sen T, Dutta A, Chatterjee A. Epigallocatechin-3-gallate (EGCG) downregulates gelatinase-B (MMP-9) by involvement of FAK/ERK/NFkappaB and AP-1 in the human breast cancer cell line MDA-MB-231. *Anticancer Drugs* 2010; 21(6): 632-44.
  23. Wei H, Wang S, Zhen L, Yang Q, Wu Z, Lei X, et al. Resveratrol attenuates the blood-brain barrier dysfunction by regulation of the MMP-9/TIMP-1 balance after cerebral ischemia reperfusion in rats. *J Mol Neurosci* 2015; 55(4): 872-9.
  24. Palmieri D, Aliakbarian B, Casazza AA, Ferrari N, Spinella G, Pane B, et al. Effects of polyphenol extract from olive pomace on anoxia-induced endothelial dysfunction. *Microvasc Res* 2012; 83(3): 281-9.
  25. Newby AC. Metalloproteinase production from macrophages-a perfect storm leading to atherosclerotic plaque rupture and myocardial infarction. *Exp Physiol* 2016; 101(11): 1327-37.
  26. Sang QX, Jin Y, Newcomer RG, Monroe SC, Fang X, Hurst DR, et al. Matrix metalloproteinase inhibitors as prospective agents for the prevention and treatment of cardiovascular and neoplastic diseases. *Curr Top Med Chem* 2006; 6(4): 289-316.

**How to cite this article:** Farrokhi E, Ghatreh-Samani K, Salehi-Vanani N, Mahmoodi A. **The effect of resveratrol on expression of matrix metalloproteinase 9 and its tissue inhibitors in vascular smooth muscle cells.** *ARYA Atheroscler* 2018; 14(4): 157-62.